

AD-A043 731

CLEMSON UNIV S C DEPT OF MATHEMATICAL SCIENCES

F/G 12/1

A NOTE ON POLYNOMIAL MATRIX FUNCTIONS OVER A FINITE FIELD.(U)

AUG 77 J V BRAWLEY

N00014-76-C-0130

UNCLASSIFIED

TR-256

NL

| OF |
AD
A043731



END
DATE
FILMED

9 -77

DDC

**DEPARTMENT
OF
MATHEMATICAL
SCIENCES**

CLEMSON UNIVERSITY
Clemson, South Carolina



A NOTE ON
POLYNOMIAL MATRIX FUNCTIONS
OVER A FINITE FIELD

BY


J. V. BRAWLEY*

DEPARTMENT OF MATHEMATICAL SCIENCES
CLEMSON UNIVERSITY


TECHNICAL REPORT #256 ✓

CONTRACT REPORT N8 ✓

*RESEARCH SUPPORTED BY
THE OFFICES OF NAVAL RESEARCH
CONTRACT N00014-76-C-0130



REPRODUCTION IN WHOLE OR PART IS PERMITTED FOR ANY PURPOSES OF
THE U.S. GOVERNMENT. DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED.



A Note on Polynomial Matrix Functions
over a Finite Field
by J.V. Brawley*

1. Let $F = GF(q)$ denote the finite field of order q , and let F_n denote the ring of $n \times n$ matrices over F . Consider an element $A(x) \in F_n[x]$; i.e.,

$$(1) \quad A(x) = A_N x^N + A_{N-1} x^{N-1} + \dots + A_1 x + A_0$$

where $A_i \in F_n$. This polynomial defines via substitution several functions from F_n to F_n . Two such functions are

$$(2) \quad B \rightarrow A_R(B) = A_N B^N + A_{N-1} B^{N-1} + \dots + A_1 B + A_0$$

and

$$(3) \quad B \rightarrow A_L(B) = B^N A_N + B^{N-1} A_{N-1} + \dots + B A_1 + A_0.$$

We call (2) and (3), respectively, the right and left polynomial functions determined by $A(x)$ with the terms right and left indicating the side on which the substituting variable is placed.

*Supported in part by ONR contract N00014-76-C-0130.

ACCESSION for		
NTIS	White Section	<input checked="" type="checkbox"/>
DDC	Buff Section	<input type="checkbox"/>
UNANNOUNCED		<input type="checkbox"/>
JUSTIFICATION		
BY		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL.	and/or SPECIAL
A		

Definition. A function $A: F_n \rightarrow F_n$ is called a right respectively left) polynomial function if there exists a polynomial $A(x) \in F_n[x]$ which represents A via the right substitution (2) (respectively (3)).

In this note we obtain unique representations for and determine the number of right (left) polynomial functions $A: F_n \rightarrow F_n$. Proofs will be given for the right functions which can be obviously modified for the left polynomial functions.

2. Recall that

$$(4) \quad L_n(x) = \prod_{i=1}^n (x^{q^i} - x)$$

is the monic polynomial of least degree in $F[x]$ satisfied by every $B \in F_n$; indeed, $L_n(x)$ is the least common multiple of all degree n polynomials in $F[x]$ [See, 2]. We define δ by

$$(5) \quad \delta = \deg L_n(x) = q^n + q^{n-1} + \dots + q.$$

THEOREM 1. Let $Z(x) = \sum_{i=0}^N Z_i x^i$ be a polynomial in $F_n[x]$ with $\deg Z(x) = N < \delta$. If $Z_r(B) = Z_N B^N + \dots + Z_1 B + Z_0 = 0$ for every $B \in F_n$, then $Z_i = 0$, $i = 0, 1, 2, \dots, N$.

Proof. Let $f(x) = x^n - a_{n-1}x^{n-1} - \dots - a_1x - a_0$ be an arbitrary polynomial of degree n in $F[x]$, and let $C \in F_n$ denote the companion matrix of $f(x)$. Dividing $Z(x)$ by $f(x)$ we obtain

$$(6) \quad Z(x) = Q(x) f(x) + R(x)$$

where $Q(x)$ and $R(x)$ are in $F_n[x]$ with

$$(7) \quad R(x) = R_{n-1}x^{n-1} + \dots + R_1x + R_0.$$

Since $f(x)$ is a scalar polynomial we may substitute an arbitrary matrix B into (6) to get $Z_r(B) = Q_r(B)f(B) + R_r(B)$. In particular, for every nonsingular $P \in GL(n, q)$ it follows from the Hamilton-Cayley theorem that

$$0 = Z_r(PCP^{-1}) = R_r(PCP^{-1}).$$

Thus $(R_r(PCP^{-1}))P = 0$ or

$$(8) \quad R_{n-1}PC^{n-1} + R_{n-2}PC^{n-1} + \dots + R_1PC + R_0P = 0$$

for every $P \in GL(n, q)$.

Now it is known [1] that each matrix $X \in F_n$ can be written as a linear combination of nonsingular matrices P_i ; i.e.,

$$X = c_1P_1 + c_2P_2 + \dots + c_tP_t, \quad c_i \in F.$$

It follows from (8) that

$$(9) \quad R_{n-1}XC^{n-1} + R_{n-2}XC^{n-1} + \dots + R_1XC + R_0X = 0$$

for every $X \in F_n$. In particular, if we take $X = E_m$ where E_m has a 1 in position $(m, 1)$ and zeros elsewhere we find through actual computation that equation (9) reduces to

$$\begin{pmatrix}
 r_{1m}^{(0)} & r_{1m}^{(1)} & \cdot & \cdot & \cdot & r_{1m}^{(n-1)} \\
 r_{2m}^{(0)} & r_{2m}^{(1)} & \cdot & \cdot & \cdot & r_{2m}^{(n-1)} \\
 \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
 r_{nm}^{(0)} & r_{nm}^{(1)} & & & & r_{nm}^{(n-1)}
 \end{pmatrix} = 0$$

where $R_k = (r_{ij}^{(k)})$. Thus column m of R_k is zero for $k = 0, 1, \dots, n-1$ and $m = 1, 2, \dots, n$; i.e., $R_k = 0$ for $k = 0, 1, \dots, n-1$. It follows from (6) that $f(x)$ divides $Z(x)$ for every monic of degree n ; hence $L_n(x)$ divides $Z(x)$. But $\deg Z(x) < \deg L_n(x)$ so $Z(x)$ must be the zero polynomial; i.e., every $Z_i = 0$ and the proof is complete.

As a corollary to Theorem 1 we have the following:

THEOREM 2. Each right polynomial function $A: F_n \rightarrow F_n$ can be represented uniquely by a polynomial $A(x) \in F_n[x]$ of degree $< \delta$ and each such polynomial represents a right polynomial function. The number of right polynomial functions is therefore $q^{n^2\delta}$.

Proof. If $A_1(x)$ and $A_2(x)$ have degree $< \delta$ and each represent the right polynomial function A then $A_1(x) - A_2(x)$ represents the zero function; hence by Theorem 1, $A_1(x) = A_2(x)$.

Finally let A be a right polynomial function and let $A(x)$

represent A. By division

$$A(x) = Q(x)L_n(x) + R(x)$$

where $R(x)$ has degree $< \delta$. Clearly, $R(x)$ represents A.

References

1. J. V. Brawley. On the ranks of basis of vector spaces of matrices. Linear Algebra and Its Applications. 3(1970), 51-55.
2. J. V. Brawley, L. Carlitz, and Jack Levine. Scalar polynomial functions on the $n \times n$ matrices over a finite field. Linear Algebra and Its Applications. 10(1975), 199-217.

Unlimited

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Department of Mathematical Sciences ✓ Clemson University Clemson, South Carolina		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3. REPORT TITLE A Note on Polynomial Matrix Functions Over a Finite Field			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) 9 Technical report			
5. AUTHOR (Last name, middle initial, first name) J. V. Brawley 14 TR-256, CR-N8			
6. REPORT DATE 8/1/77 91/1 Aug 77		7a. TOTAL NO. OF PAGES 5	7b. NO. OF REFS 2
8a. CONTRACT OR GRANT NO. N00014-76-C-0130 ✓		9a. ORIGINATOR'S REPORT NUMBER(S) N8 ✓	
8b. PROJECT NO.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) 92 8p.	
10. DISTRIBUTION STATEMENT Unlimited			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY 407 183	
13. ABSTRACT Let $F = GF(q)$ denote the finite field of order q , and let F_n denote the ring of $n \times n$ matrices over F . Each matrix polynomial $A(x) = A_N x^N + \dots + A_1 x + A_0$ in $F_n[x]$ defines via substitution several functions from F_n to F_n . Two such functions, called respectively, the right and left polynomial functions determined by $A(x)$ are $B \rightarrow A_r(B) = A_N B^n + \dots + A_1 B + A_0$ $B \rightarrow A_L(B) = B^n A_N + \dots + B A_1 + A_0$ A function $A: F_n \rightarrow F_n$ is called a right (left) polynomial function if there exists $A(x) \in F_n[x]$ which represents A via the right (left) substitution $B \rightarrow A_r(B)$ ($B \rightarrow A_L(B)$). This paper obtains a unique representation for and determines the number of right (left) polynomial functions $A: F_n \rightarrow F_n$.			